

Crush Injury of Upper Extremity Leading to Bywater's Syndrome: A Case Report

Anis Puspita Utami¹, Rahmadia Kusumamardhika², Agus Maruli³

¹⁻³Department of Orthopaedics,
University of Brawijaya, Indonesia

Authorship and contribution Declaration:

Each author of this article fulfilled ALL 03 Criteria of Authorship:

1. Conception and design of or acquisition of data or analysis and interpretation of data.
2. Drafting the manuscript or revising it critically for important intellectual content.
3. Final approval of the version for publication.
4. All authors agree to be responsible for all aspects of their research work

Corresponding author:

Anis Puspita Utami

E-mail:

anispuspita.utami@gmail.com

ABSTRACT

A male came to the ER with wounds on his left arm after a vehicle incident. He was run over by the truck's wheels on his left upper extremity until his whole arms were crushed. He had a massive crush injury on his whole left arm, no movement and pulse were detected. MESS Scoring resulted in score of 11, thus an indication for amputation. Blood examination showed increase of ureum and creatinine without previous kidney disease history. The patient was diagnosed with a crush injury of the upper extremity and Bywater's Syndrome based on kidney involvement. He was consulted to the orthopedic department and was given fluid resuscitation, injections of analgesic, antibiotics, and Human Tetanus Immunoglobulin (HTIG). Proximal humeral amputation was performed. A second surgery later was performed to debride the wound due to contaminated necrotic tissues.

Keywords: Accidents, Traffic, Accident, Bone, Fracture, Injury, Motorbike, Pattern.

This article may
be cited as:

Utami, A.P., Kusumamardhika, R., Maruli, A. Crush Injury of Upper Extremity Leading to Bywater's Syndrome: A Case Report. *J. Pak. Orthop. Assoc.* 2024; Vol. 36 (01):60-62.

INTRODUCTION

Crush injury can lead to many complications, one such uncommon consequence is known as Bywater's Syndrome.¹ It is characterized by the sudden and excessive destruction of red blood cells (hemolysis) due to mechanical trauma.² It is a relatively rare condition, Mangled Extremity Severity Score (MESS) should be performed to patients with crush injury as diagnostic tools and to determine if the patient needs amputation.³

CASE REPORT

A 57-year-old male came to the ER with a massive wound on his left arm after a motor vehicle incident on the street. Patient was riding his bike and then collided with a truck. Patient got run over by the truck's wheels on his left upper extremity until his whole arms were crushed.

Vital signs showed blood pressure of 85/51mmHg, body temperature of 36.5°C, heart rate of 128 bpm, respiratory rate of 24 times per minute. Physical examination showed a massive crush injury on his whole left arm, no movement and pulse detected. MESS Scoring was performed and the

patient's score is 11, thus an indication for amputation. The left humerus x-ray was performed even though the crush injury was already visible. The x-ray showed one-third mid left humeral fracture with poor alignment. Blood examination showed lowered hemoglobin of 10.4 g/dl, elevated leukocytes of 34.720/uL, urea and creatinine of 59 mg/dL and 2.12 mg/dL without previous kidney disease history. The patient was diagnosed with a crush injury of the upper extremity and Bywater's Syndrome due to kidney involvement. The patient was consulted to the orthopedic department and was given 2 liter of ringer lactate fluid, ketorolac injection, ceftriaxone injection, and Human Tetanus Immunoglobulin (HTIG) and proximal humerus amputation was performed. Three days later, he complained about darkened skin around his amputated arm, followed by slight blood and pus. A second surgery was performed to debride the wound due to contaminated necrotic tissues.

Five days after second surgery, the patient showed improvement and there were no signs of infection. The patient was discharged from hospital and continue the treatment to evaluate post amputation condition and systemic symptoms.



Fig. 1: Crushed left arm after injury




Fig. 2: Upper left arm x-ray



Fig. 3: Necrotizing tissue after amputation surgery



Fig. 4: Debridement surgery



Fig. 5: Post debridement surgery

DISCUSSION

A crush injury is characterized by the extensive damage of a large muscle mass. Bywater's syndrome or crush syndrome is a crush injury with systemic manifestations.⁴ Systemic symptoms arise from traumatic rhabdomyolysis, which occurs when there is muscle reperfusion injury following the release of compressive forces on the tissues.⁵ The patient showed kidney dysfunction as in elevated creatinine and urea level upon trauma without prior kidney failure history. Upon arrival, vital signs showed hypotension, tachycardia, and tachypnea, which might be early signs of hypovolemic shock due to

massive blood loss. Early resuscitation and treatment were done to prevent further complication. According to literature, early untreated Bywater's syndrome death is caused by hypovolemic shock and hyperkalemia due to kidney dysfunction, and late untreated death is caused by prolong renal failure, coagulopathy, hemorrhage, and sepsis.⁶ Further treatment is determined by using a MESS score to decide whether the patient needs an amputation. MESS score includes skeletal/soft tissue injury, limb ischemia, shock, and age as variables.⁷ Score of 7 or above is highly predictive of amputations. As this patient scored 11, amputation surgery was performed. According to literature, it is reported 13-40% of infection happen in major limb amputation. The risk factors varies from ischemia, pre-existing limb ulcers, patient co-morbidities, and contamination of the wound.⁸ In this patient, contamination of the wound is verdict to become the source of infection. The second surgery was performed to debride the contaminated wound. After a total of six days of treatment and two surgeries, the patient showed improvement after the second surgery. Further treatment were still needed to evaluate the wound and systemic signs in kidney involvement. Post-operative pain and psychological factors are also needed to be issued. We would recommend psychological assessment, kidney function evaluation, and wound hygiene for the next treatment. It is an important case to be reported as early recognition and treatment might result in a notably outcome for the patient.

CONCLUSION

Urgent and effective medical care is required to reduce the risk of cardiac arrhythmia, kidney damage and death. Decisions may need to be made quickly as postponed therapy might result in worse prognosis. Patients with crush injury present some of

the greatest challenges in field medicine, and may need a physician's attention on the site of their injury. Appropriate physiological preparation of the injured is mandatory.

Conflict of Interest: None

Grants/Funding: None

REFERENCES

1. Long B, Liang SY, Gottlieb M. Crush injury and syndrome: a review for emergency clinicians. *The American Journal of Emergency Medicine*. 2023 Apr 25.
2. Yücesoy M, Dünya CP. Crush Syndrome and Nursing Care Management. *CURARE Journal of Nursing*. 2023 Jul 7(2):49-55.
3. Mahmoud AA, Elbahat MA. Validity of Mangled Extremity Score System to Predict Limb Salvage on Cases of Traumatic Popliteal Artery Injuries: A Case Series and Review of the Literature. *Journal of Endovascular Resuscitation and Trauma Management*. 2023 Jul 4.
4. Xu P, Wang F, Zhou XL, Li L, Xiong D, Yong YQ, Zhao Y, Jiang WX. Systemic inflammatory response and multiple organ dysfunctions following crush injury: a new experimental model in rabbits. *Inflammation*. 2018 Feb;41:240-8.
5. Dhatt SS, Neradi D. Managing Crush Injuries on Arrival. *InAcute Trauma Care in Developing Countries* 2022 Sep 2 (pp. 140-146). CRC Press.
6. Loja MN, Sammann A, DuBose J, Li CS, Liu Y, Savage S, Scalea T, Holcomb JB, Rasmussen TE, Knudson MM, AAST PROOVIT Study Group. The mangled extremity score and amputation: time for a revision. *The journal of trauma and acute care surgery*. 2017 Mar;82(3):518.
7. Vasei N, Jahangiri K. A case report of crush syndrome and delay in invasive therapeutic intervention: is the priority to save the life of the patient or to maintain the limb. *Journal of Safety Promotion and Injury Prevention*. 2020;7(3).
8. Sadat U, Chaudhuri A, Hayes PD, Gaunt ME, Boyle JR, Varty K. Five day antibiotic prophylaxis for major lower limb amputation reduces wound infection rates and the length of in-hospital stay. *European Journal of Vascular and Endovascular Surgery*. 2008 Jan 1;35(1):75-8.